


WHAT IS CLAIMED IS:

1. A metal coordination compound represented by the following formula (1):



5 wherein M denotes Ir, Pt, Ph or Pd; L denotes a bidentate ligand; L' denotes a bidentate ligand different from L; m is an integer of 1, 2 or 3; and n is an integer of 0, 1 or 2 with the proviso that the sum of m and n is 2 or 3,

10 the partial structure  $M L_m$  being represented by a formula (2) or a formula (3) shown below, and the partial structure  $M L'_n$  being represented by a formula (4) or a formula (5) shown below:



wherein CyN1, CyN2 and CyN3 independently denote a substituted or unsubstituted cyclic group containing a nitrogen atom connected to M; CyN4 denotes a cyclic group containing 8-quinoline or its derivative having a nitrogen atom connected to M; CyC1, CyC2 and CyC3 independently denote a substituted or unsubstituted cyclic group containing a carbon atom connected to M,

25 each of substituents for CyN1, CyN2, CyN3, CyC1, CyC2 and CyC3 being selected from the group consisting of a halogen atom; cyano group; nitro

group; a trialkylsilyl group containing three linear  
or branched alkyl groups each independently having 1 -  
8 carbon atoms; a linear or branched alkyl group  
having 1 - 20 carbon atoms capable of including one  
5 or at least two non-neighboring methylene groups which  
can be replaced with -O-, -S-, -CO-, -CO-O-, -O-CO-,  
-CH=CH- or -C≡C- and capable of including a hydrogen  
atom which can be replaced with a fluorine atom; and  
an aromatic ring group capable of having a substituent  
10 selected from the group consisting of a halogen atom;  
cyano group; nitro group; and a linear or branched  
alkyl group having 1 - 20 carbon atoms capable of  
including one or at least two non-neighboring  
methylene groups which can be replaced with -O-, -S-,  
15 -CO-, -CO-O-, -O-CO-, -CH=CH- or -C≡C- and capable of  
including a hydrogen atom which can be replaced with a  
fluorine atom,

CyN1 and CyC1 being connected via a covalent  
group containing X which is represented by -O-,  
20 -S-, -CO-, -C(R1)(R2)- or -NR- where R1, R2 and R  
independently denote a hydrogen atom, a halogen  
atom, an alkyl group, an alkyl group substituted  
with a halogen atom, a phenyl group or a naphthyl  
group, and

25 CyN2 and CyC2, and CyN3 and CyC3 being  
independently connected via a covalent bond,

with the proviso that the metal coordination


00995608-12904  
FOGAT-B055650

compound is represented by the formula (2) when n is 0.

2. A compound according to Claim 1, wherein the partial structure MLM is represented by the formula (2).

3. A compound according to Claim 2, wherein M is Ir.

4. A compound according to Claim 2, wherein the metal coordination compound has another partial structure represented by the following formula (6):



wherein M denotes Ir, Pt, Ph or Pd; p is 1; and E and G independently denote a linear or branched alkyl group having 1 - 20 carbon atom capable of including a hydrogen atom which can be replaced with a fluorine atom, or an aromatic ring group capable of having a substituent selected from the group consisting of a halogen atom; cyano group; nitro group; a trialkylsilyl group containing three linear or branched alkyl groups each independently having 1 - 8 carbon atoms; and a linear or branched alkyl group having 1 - 20 carbon atoms capable of including one or

at least two non-neighboring methylene groups which can be replaced with -O-, -S-, -CO-, -CO-O-, -O-CO-, -CH=CH- or -C≡C- and capable of including a hydrogen atom which can be replaced with a fluorine atom.

5

5. A compound according to Claim 1, which exhibits a phosphorescence at the time of energy transition from an excited state to a ground state.

10

6. A compound according to Claim 1, wherein one of the ligands L and L' is a luminescent ligand and the other ligand is a carrier transport ligand.

15

7. A compound according to Claim 1, wherein at least one of the ligands L and L' is in a metal to ligand charge transfer excited state.

20

8. A compound according to Claim 1, wherein the ligands L and L' includes a first ligand capable of providing a first maximum luminescence wavelength  $\lambda_1$  based on an excited state thereof and a second ligand capable of providing a second maximum luminescence wavelength  $\lambda_2$  shorter than  $\lambda_1$ , the number of the first ligand providing  $\lambda_1$  being smaller than that of the second ligand providing  $\lambda_2$ .

25

9. A compound according to Claim 1, wherein the

0095638 1 2901  
10521 3095550

ligands L and L' includes a stronger luminescent ligand and a weaker luminescent ligand, the number of the stronger luminescent ligand is smaller than that of the weaker luminescent ligand.

5

10. An organic luminescence device, comprising: a substrate, a pair of electrodes disposed on the substrate, and a luminescence function layer disposed between the pair of electrodes comprising at least one species of an organic compound,

10

wherein the organic compound comprises a metal coordination compound represented by the following formula (1):




15

wherein M denotes Ir, Pt, Ph or Pd; L denotes a bidentate ligand; L' denotes a bidentate ligand different from L; m is an integer of 1, 2 or 3; and n is an integer of 0, 1 or 2 with the proviso that the sum of m and n is 2 or 3,


20

the partial structure  $M L_m$  being represented by a formula (2) or a formula (3) shown below, and the partial structure  $M L'_n$  being represented by a formula (4) or a formula (5) shown below:


25




(2),



(3),



(4),



(5),

wherein CyN1, CyN2 and CyN3 independently denote a substituted or unsubstituted cyclic group containing a nitrogen atom connected to M; CyN4 denotes a cyclic group containing 8-quinoline or its derivative having a nitrogen atom connected to M; CyC1, CyC2 and CyC3 independently denote a substituted or unsubstituted cyclic group containing a carbon atom connected to M, each of substituents for CyN1, CyN2, CyN3, CyC1, CyC2 and CyC3 being selected from the group consisting of a halogen atom; cyano group; nitro group; a trialkylsilyl group containing three linear or branched alkyl groups each independently having 1 - 8 carbon atoms; a linear or branched alkyl group having 1 - 20 carbon atoms capable of including one or at least two non-neighboring methylene groups which can be replaced with -O-, -S-, -CO-, -CO-O-, -O-CO-, -CH=CH- or -C≡C- and capable of including a hydrogen atom which can be replaced with a fluorine atom; and an aromatic ring group capable of having a substituent selected from the group consisting of a halogen atom; cyano group; nitro group; and a linear or branched alkyl group having 1 - 20 carbon atoms capable of including one or at least two non-neighboring methylene groups which can be replaced with -O-, -S-, -CO-, -CO-O-, -O-CO-, -CH=CH- or -C≡C- and capable of including a hydrogen atom which can be replaced with a fluorine atom,

00995508 112001

CyN1 and CyC1 being connected via a covalent group containing X which is represented by -O-, -S-, -CO-, -C(R1)(R2)- or -NR- where R1, R2 and R independently denote a hydrogen atom, a halogen atom, an alkyl group, an alkyl group substituted with a halogen atom, a phenyl group or a naphthyl group, and

CyN2 and CyC2, and CyN3 and CyC3 being independently connected via a covalent bond,

with the proviso that the metal coordination compound is represented by the formula (2) when n is 0.

11. A device according to Claim 10, wherein the partial structure MLm is represented by the formula (2).

12. A device according to Claim 11, wherein M is Ir.

13. A device according to Claim 10, wherein a voltage is applied between the pair of electrodes to cause phosphorescence from the luminescence function layer.

14. An image display device, comprising: an organic luminescence device according to Claim 10 and

means for supplying electrical signals to the organic luminescence device.

5

10

15

20

25